

ISSN 2224-4980



**International
Journal of Ecosystems and Ecology Science
(IJEES)**

<https://doi.org/10.31407/ijeess>



Essays on Ecosystems and Environmental Research

Volume 13/2, 2023

February 2023- June 2023

<http://ijeess.net/>

**International
Journal of Ecosystems and Ecology Science
(IJEES)**

<https://doi.org/10.31407/ijeess>





**International Journal of Ecosystems and Ecology Science
(IJEES), ISSN: 2224-4980**

Volume 13, Issue 2, February 2023 - June 2023, (Serial Number 50)



Editor in Chief

Hysen Mankolli, Professor, Ecologist, Plainfield, Illinois, USA; Health and Environment Association, Tirana Albania.
Earth System Science Interdisciplinary Center (ESSIC), University of Maryland College Park, USA, Consultant Scientist.

Deputy-Editor-in-Chief

Sukru Dursun, Professor, Konya Technical University, Environmental Engineering Dep., Konya-Turkey.
Massimo Zuchetti, Professor, MIT, Massachusetts Institute of Technology, Cambridge MA, USA.
Cezar Kongoli, Professor, University of Maryland College Park, Earth System Science Interdisciplinary Center (ESSIC), USA.
Selim Dogan, Professor, Konya Technical University, Environmental Engineering Dep., Konya-Turkey.

International Scientific Editorial Board

Sukru Dursun, Professor, Konya Technical University, Environmental Engineering Dep., Konya, Turkey;
Massimo Zuchetti, Professor, MIT, Massachusetts Institute of Technology, Cambridge MA, USA;
Veselin Alexandrov, Professor, National Institute of Meteorology and Hydrology, Bulgaria;
Nicola Senesi, Professor, University of Bari, Bari, Italy;
Wander Michelle M, Professor, University of Illinois at Urbana-Champaign, Urbana, Urbana, IL 61801, USA;
Vladimir Pesic, Professor, University of Montenegro, Department of Biology, Podgorica Montenegro;
Ibraim Dincer, Professor, University of Ontario, Faculty of Engineering and Applied Science, Ontario, Canada;
Hysen Mankolli, Professor, Ecologist, Plainfield, Illinois, USA;
Lyudmyla Symochko, Prof. Assoc. Dr. Uzhhorod National University, Faculty of Biology, Uzhhorod, Ukraine;
Zacharoula S. Andreopoulou, Professor, Aristotle University of Thessaloniki, Thessaloniki, Greece;
Mariana Golumbeanu, PhD, National Institute for Marine Research and Development "Grigore Antipa", Constanta, Romania;
Antonis K. Kokkinakis, Professor, Aristotle University of Thessaloniki, Thessaloniki, Greece;
Olivia Cioboiu, PhD, The Oltenia Museum Craiova, Popa Șapcă str. No 8, Rumania;
Nasser Modirshahla, Professor, Islamic Azad University, Department of Applied Chemistry, Iran;
Nina Liogchii, Professor, Institut of Ecology and Geografy, Republic of Moldova;
Trajce Talevski, Professor, Hydrobiological Institute, Naum Ohidski 50, 6000 Ohrid - R.Macedonia;
Alexander V. Derunkov, Professor, Dep. of Entomology, National Museum of Natural History, Washington DC, USA;
Can Ozgur Colpan, Professor, Ryerson University, Canada;
Muhammad Ashraf, Professor, University of Agriculture, Faculty of Sciences, Faisalabad 38040, Pakistan;
Roberto Mancinelli, Professor, University of Tuscia, Viterbo, Italy;
Bahrive Gulgun, Professor, Ege University, Izmir, Turkey;

Insaf Mekki, PhD, National Research Institute for Rural Engineering, Water, and Forestry (I.N.R.G.R.E.F), Ariana, Tunisia;
Hany Gaber El Shaer, PhD, IUCN Centre for Mediterranean Cooperation / Global Marine, Egypt;
Ertugrul Esmeray, Ph.D, Karabuk University, Engineering Faculty, Environmental Engineering Department Turkey;
Violeta Vidaček-Hainš, PhD, Ass. Professor, University of Zagreb, Croatia;
Cezar Kongoli, Professor, University of Maryland College Park, Earth System Science Interdisciplinary Center (ESSIC), USA;
Justas Kažys, PhD, Assoc. Professor, Vilnius University, Department of Hydrology and Climatology, Vilnius –Lithuania;
Selim Dogan, PhD, Assoc. Professor Selcuk University, Environmental Engineering Department, Konya Turkey;
Hisham Mostafa Alidrisi, PhD, Assoc. Professor, King Abdulaziz University, Engineering College, Jeddah Saudi, Arabia;
Khalid A. A. Al-Ghamdi, PhD, Assoc. Professor, King Abdulaziz University, Engineering College, Jeddah Saudi, Arabia;
Osman Taylan, PhD, Assoc. Professor, King Abdulaziz University, Engineering College, Jeddah Saudi, Arabia;
Robert J. Kuligowski, PhD, University of Maryland College P, Earth System Science Interdisciplinary Center (ESSIC), USA;
Mariia M. Fedoriak, Professor, Chernivtsi National University, Dep. of Ecology and Biomonitoring, Chernivtsi, Ukraine;
Narayan Ramappa Birasal, Associate Professor, KLE Society's G H College, Zoology Department, , Karnataka state, India;
Rana P. Singh, Professor, Babasaheb Bhimrao Ambedkar University, Dep. of Environmental Science, U.P., India;
Hayati AKMAN, Dr., Selcuk University, Seed Department, 42430 Konya, Turkey;
Fatma ÇANKA KILIÇ, Professor, Kocaeli University, Technology Faculty, Energy Systems Engineering Dep., Kocaeli, Turkey;
Ferim Gashi, Dr., University of Pristina, Department of Geography, Faculty of Mathematical-Natural Sciences, Kosovo;
Altin Dorri, Professor Assoc., Polytechnic University of Tirana, Tirana, Albania;
Alexander P. Sizykh, Professor, Russian Academy of Sciences, Siberian Institute, Irkutsk, Lermonyova, Russia;
Zeynep EREN, Professor, Ataturk University, Environmental Engineering Department, Erzurum, Turkey;
Sheida Korjani, PhD, Department of Architecture, Tabriz Branch, Islamic Azad University, Tabriz, Iran;
Maan Maarroof, PhD, Assistant Professor, University of Mosul, College of Veterinary Medicine, Iraq;
Koula Doukani, Professor, University of Ibn Khaldoun, Zaaroura, Tiaret, Algeria;
Svetla Petkova Gateva, PhD, Institute of Biodiversity and Ecosystem Research, Bulgarian Acad. of Sciences, Sofia, Bulgaria;
Rrahim Sejdiu, Professor Assistant, PhD, University of Applied Sciences-Ferizaj, Kosovo;
Mouloud BOUHOUBOU, Dr., ENS Assia Djebar - Constantine, Algeria;
Dinh Tran Ngoc Huy, Dr., Banking University HCMC, Ho Chi Minh city Vietnam-International University of Japan, Japan;
Admir Jançe, Professor Assoc., European University of Tirana, Tirana, Albania;
Nataliia P. Kovalenko, Dr., Institute of Plant Physiology and Genetics of NAS of Ukraine, Kyiv, Ukraine;
Leila Soudani, Dr. University of Tiaret, Faculty of Nature and Life Sciences, Tiaret Algeria;

Publication Information:

International journal of ecosystems and ecology science (IJEES), ISSN 2224-4980, publishing original articles, reviews and short communications of high scientific standard on ecosystems and science ecology. By: Hysen Mankolli, Professor, Ecologist, Plainfield, Illinois, USA; Health and Environment Association, Tirana, Albania;

ISSN International Centre, *Bibliographic Data Section*, PARIS, FRANCE;

Bib - ID 68337; ISO standard 3297;

Key title: International journal of ecosystems and ecology science

Abbreviated key title: *Int. j. ecosyst. ecol. sci.*

ISSN 2224-4980; <http://ijees.net/>

Frequency: four times a year. IJEES Electronic Journal Publication: Plainfield, Illinois, USA;

DOI prefix: 10.31407, <https://doi.org/10.31407/ijees>

IJEES doi crossref

<https://doi.org/10.31407/ijees>

EBSCO's library products, USA

<http://www.ebscohost.com/for-publishers/for-publishers>

Index Copernicus, ICV= 2020: **100.00**; ICV= 2019: **100.00**; ICV= 2018: **100.00**; ICV= 2017: **98.95**;

<https://journals.indexcopernicus.com/search/journal/issue?issueId=197247&journalId=31407>

CiteFactor, **IF 2019-2020=1.40**

<https://www.citefactor.org/impact-factor/impact-factor-of-journal-International-Journal-of-Ecosystems-and-Ecology-Science-IJEES.php>

Scientific Journal Impact factor, **SJIF: 2021: 6.121; 2020: 7.167; 2019: 7.64**

<http://sjifactor.com/passport.php?id=17557>

SCILIT

<https://app.scilit.net/publications?source-id=23651&type=journal-article>

Academic Resource Index Research Bible:

<http://journalseeker.researchbib.com/view/issn/2224-4980>

Google scholar Cited, **h-index: 10; Citations 510**;

<https://scholar.google.com/citations?user=W89N8qgAAAAJ&hl=en>

MIAR, **ICDS 2021: 9.5**;

<http://miar.ub.edu/issn/2224-4980>

Web of Science Master List 2018-2022

<https://rdmusm.files.wordpress.com/2020/01/journal-list-emerging-sources-citation-index-web-of-science-update-on-23-jan-2020.pdf>

UOI license: <http://u-o-i.org/1.01/ijees>; <http://www.u-o-i.org/index.php/SearchLicense>



DPI Digital Library USA and assign unique digital no. DPI: 16.10047.IJEES

U.S. National Library of Medicine.

NLM ID: [101726509](https://www.ncbi.nlm.nih.gov/nlmcatalog/101726509) [Serial]; Other ID: (OCOLC)879805056;

<https://www.ncbi.nlm.nih.gov/nlmcatalog/101726509>

This journal has the status of an international journal.

Aims and Scope:

The goals of the International Journal of Ecosystems and Ecology Science (IJEES), ISSN 2224-4980, are to bring together researchers and scientists with interests in the quality of the ecosystems research results, theories, technologies, systems, tools, applications, the work in progress and experiences on ecosystems used. The main topics of interest are:

- ✓ Ecosystems
- ✓ Agro ecosystems
- ✓ Forest ecosystems
- ✓ Ecology
- ✓ Plant Ecology
- ✓ Animal Ecology
- ✓ Human Ecology
- ✓ Lakes and Rivers ecosystems
- ✓ Applied Biology
- ✓ Applied Ecology
- ✓ Applied Chemistry
- ✓ Biodiversity
- ✓ Energy
- ✓ Geology
- ✓ Bioaccumulation
- ✓ Desalination
- ✓ Water
- ✓ Soil
- ✓ Air pollution
- ✓ Climate Change
- ✓ Ecosystem restoration
- ✓ Environment toxicology
- ✓ Environment protection
- ✓ Environmental radioactivity
- ✓ Environmental legislation
- ✓ Environmental management
- ✓ Environmental education.
- ✓ Green technology
- ✓ Socioeconomic aspects in Ecosystems
- ✓ Agro tourism and National Park
- ✓ Health Care
- ✓ Food security
- ✓ Rights and Environmental Laws
- ✓ Environment Engineering
- ✓ Environment Architecture
- ✓ Eco philosophy Issue

Abstracted / Indexed / Impact Factor in:

International journal of ecosystems and ecology science (IJEES), ISSN 2224-4980,

IJEES: <http://ijeess.net/>

IJEES have an international status and has been indexed, abstracted, and have impact factors:

ISSN Center responsible of the record: CIEPS - ISSN

<https://portal.issn.org/resource/ISSN/2224-4980>

IJEES doi crossref

<https://doi.org/10.31407/ijeess>

IJEES, Volume 13/2, 2023

[https://ijeess.net/journal-89-International-Journal-of-Ecosystems-and-Ecology-Science-\(IJEES\)-Volume-13-2,-2023.html](https://ijeess.net/journal-89-International-Journal-of-Ecosystems-and-Ecology-Science-(IJEES)-Volume-13-2,-2023.html)

EBSCO's library products,USA

<http://www.ebscohost.com/for-publishers/for-publishers>

Science gate Cited

<https://www.sciencegate.app/source/315335/top-cited>

Index Copernicus, ICV= 2020: 100.00; ICV= 2019: 100.00; ICV= 2018: 100.00; ICV= 2017: 98.95;

<https://journals.indexcopernicus.com/search/journal/issue?issueId=197247&journalId=31407>

CiteFactor, IF 2019-2020=1.40

<https://www.citefactor.org/impact-factor/impact-factor-of-journal-International-Journal-of-Ecosystems-and-Ecology-Science-IJEES.php>

Scientific Journal Impact factor, SJIF: 2021: 6.121; 2020: 7.167; 2019: 7.64

<http://sjifactor.com/passport.php?id=17557>

SCILIT

<https://app.scilit.net/publications?source-id=23651&type=journal-article>

Academic Resource Index Research Bible:

<http://journalseeker.researchbib.com/view/issn/2224-4980>

Google scholar Cited, h-index: 10; Citations 510;

<https://scholar.google.com/citations?user=W89N8qgAAAAJ&hl=en>

MIAR, ICDS 2021: 9.5;

<http://miar.ub.edu/issn/2224-4980>

exaly

<https://exaly.com/journal/102491/international-journal-of-ecosystems-and-ecology-science-ijeess/impact-factor>

Web of Science Master List 2018-2022

<https://rdmusm.files.wordpress.com/2020/01/journal-list-emerging-sources-citation-index-web-of-science-update-on-23-jan-2020.pdf>

WorldCat

https://www.worldcat.org/search?qt=affiliate_wc_org_all&ai=Directory_ashok.drji%252540gmail.com&fq=&q=2224-4980&wcbtm2w=Go

US. National Library of Medicine, NLM ID: 101726509 [Serial]; Other ID: (OCOLC)879805056;

<https://www.ncbi.nlm.nih.gov/nlmcatalog/101726509>

CAS Chemical Abstracts Service – CAS database

<http://cassi.cas.org/search.jsp>

Environment-and-Nature/Biodiversity/Ecosystem.

<http://www.efita.org/Environment-and-Nature/Biodiversity/Ecosystem-Biodiversity/IJEES-details-28892.html>

NewJour : Electronic Journal&Newsletters

<http://gulib.georgetown.edu/newjour/nj2/index.html>

Environmental expert

<http://www.environmental-expert.com/magazines/?keyword=ecosystem+science>

Currently Indexing at Pro Quest

http://www.proquest.com/en-US/catalogs/databases/detail/periodicals_index.shtml

CABI

<http://www.cabi.org/default.aspx?page=1016&site=170&pid=40&xslttab=2&newtitlesonly=1&search=>

Academia.edu

http://www.academia.edu/3834422/2013_Networking_approach_to_the_sustainable_development_of_European_Regions_bordering_major_rivers

Manuscripts and correspondence are invited for publication

*Dear Professors, PhD, researchers and scholars of the field of ecosystems and related areas to, such as environmental engineering, agriculture, bio-chemistry, soil, water and air pollution, geology, forestry, urban ecology, resource economics, etc., at the moment, I declare the sending of scientific journal articles to International Journal of Ecosystems and Ecology Science (IJEES) is open. Sending of items can only be done electronically, to the following address: e-mail: editorijeess@gmail.com; 2011ijeese@gmail.com; mjedisishendeti@yahoo.com; The confirmation of receiving these items will be announced within the period of one to three weeks after sending. Whether the article is accepted or not, will be announced within a period of two to three weeks. Only original articles will be accepted, validated, and not to have or be published in other journals. Publication lasts from two to six months. All Rights Reserved Copyright by © IJEES (the first or correspondent author have responsible for the accuracy of the article). Submission guidelines and Web Submission System are available at: <https://www.ijeess.net/page-6-Guide-for-authors.html>;
Note: If the paper is published in another journal or if it includes false or inaccurate claims, it will be rejected or removed automatically as that constitutes a violation. Zero tolerance!*

Editorial Office:

*IJEES Electronic Journal Publication: Plainfield, Illinois, USA; E-mail: editorijeess@gmail.com;
Health and Environment Association, Tirana, Albania; E-mail: mjedisishendeti@yahoo.com;*

**International
Journal of Ecosystems and Ecology Science
(IJEES)**

DOI: <https://doi.org/10.31407/ijeess>

Volume 13/2, 2023

DOI: <https://doi.org/10.31407/ijeess13.2>

Table of contents:

Oleksandr Tkachuk*, Nadiya Viter, Snizhana Pankova, Olga Titarenko, Lyudmila Yakovets, THE CURRENT ENVIRONMENTAL STATE OF THE FIELD PROTECTIVE FOREST BELTS OF THE FOREST STEPPE OF UKRAINE, page 1-6;

DOI: <https://doi.org/10.31407/ijeess13.201>

Mohammed Balubaid¹, Waqar Ahmad Gulzar¹, Hani Aburas¹, Osman Taylan¹, Abdulaziz S. Alkabaa¹, Omer A. Bafail¹, Anas A. Makki², Ammar Y. Alqahtani¹, Hassan M. Alidrisi¹, Basil Omar Al-Sasi¹, Shoukath Ali Karuvatt³, Hisham Alidrisi^{1*}, MONITORING THE PERFORMANCE OF AGRICULTURAL AND FOOD SECTOR COMPANIES USING DEA, page 9-24;

DOI: <https://doi.org/10.31407/ijeess13.202>

DOUKANI Koula^{12*}, BOUHENNI Hasna¹, BOUKIRAT Dyhia³, HADJARAB Mohamed Hocine¹, MOKHTAR Asma¹, ZAHY Racha¹, EPIDEMIOLOGICAL STUDY OF GASTRODUODENAL DISEASES (*HELICOBACTER PYLORI*) IN THE REGION OF TIARET, ALGERIA, page 25-32;

DOI: <https://doi.org/10.31407/ijeess13.203>

Eugena Tomini^{1*}, Adela Vasili¹, Elona Kureta¹, Artan Simaku¹, Silvia Bino¹, COVID-19: A PANDEMIC EXPERIENCE FOR THE COOPERATION OF THE SCIENTIFIC COMMUNITY WORLDWIDE, page 33-36;

DOI: <https://doi.org/10.31407/ijeess13.204>

Fatma Köprülü^{1*}, Osama Abduljalil Mohammad Hamad^{2*}, Aşkın Kiraz^{3*}, PERCEPTIONS OF LIBYANS TOWARDS CLIMATE CHANGE, page 37-44;

DOI: <https://doi.org/10.31407/ijeess13.205>

Nevila Bushati^{1,2,*}, Anila Neziri², Arlind Mustafa³, PHYSICO-CHEMICAL AND BACTERIOLOGICAL EVALUATION FROM GROUNDWATERS OF PUKA CITY, page 45-50;

DOI: <https://doi.org/10.31407/ijeess13.206>

Inna Honcharuk, Oleksandr Tkachuk*, Olexandr Mazur, Ruslan Kravets, Olena Mazur, Oleksiy Alieksieiev, Zabarna Tetyana, Lina Bronnikova, MANIFESTATION OF ECOLOGICAL-ADAPTIVE PROPERTIES OF SOYBEAN VARIETIES DEPENDING ON SOIL-CLIMATIC CONDITIONS, page 51-60;

DOI: <https://doi.org/10.31407/ijeess13.207>

Mohammed Alamoudi^{1*}, SAFETY CLIMATE DIMENSIONS AS PREDICTORS FOR WORKER'S SATISFACTION REGARDING SAFETY LEVEL, page 61-70;

DOI: <https://doi.org/10.31407/ijeess13.208>

Serhii Razanov^{1,2*}, Olha Koruniak³, Andrii Dydiv¹, Tetiana Holubieva⁴, Lyudmyla Symochko^{5,6}, Volodymyr Balkovskyy¹, Oleksiy Alekseev², Oksana Vradii², Halina Ohorodnichuk², Mikhaylo Polishchuk², Oleh Kolisnyk², Oleksandr Mazur², Olena Mazur², RADIOPROTECTIVE AND SORPTION PROPERTIES OF BEESPINE, page 71-76;

DOI: <https://doi.org/10.31407/ijeess13.209>

Victor Mazur¹, Hanna Pantsyreva^{1*}, Oleh Ovcharuk², Kateryna Mazur¹, Ruslan Myalkovsky³, Oleg Tkach³, Serhii Verholiuk¹, FEATURES OF THE FUNCTIONING OF THE ASSIMILATION APPARATUS OF LUPINUS ALBUS DEPENDING ON THE USE OF ECO-BIOLOGICAL PREPARATIONS, page 77-84;

DOI: <https://doi.org/10.31407/ijeess13.210>

Furkan TÜRK^{1,2}, Hayriye Nur NAYAN², Selim DOĞAN^{3*}, Ülkü Sultan KESKİN², THE CARBON COST OF THE BUILDINGS DESTROYED IN YALOVA IN THE 1999 MARMARA EARTHQUAKE, page 85-90;

DOI: <https://doi.org/10.31407/ijeess13.211>

Alla Levishko¹, Iryna Gumeniuk^{1*}, Viktoriya Tsvigun¹, Svitlana Mazur¹, Yeuheniia Tkach¹, Olena Demyanyuk¹, Tetiana Kovalenko², THE INFLUENCE OF FIRE CULTIVATION ON THE SOIL MICROORGANISMS, page 91-96;

DOI: <https://doi.org/10.31407/ijeess13.212>

Oleg Tkach¹, Viktor Dubik¹, Oleh Ovcharuk², Lyudmila Mikhaylova¹, Hanna Pantsyreva^{3*}, Dariia Vilchynska¹, Sergii Slobodian¹, Oleg Gorbovy¹, TECHNOLOGICAL CHARACTERISTICS AND POTENTIAL OF LANDFILL BIOGAS FOR ELECTRICITY GENERATION, page 97-108;

DOI: <https://doi.org/10.31407/ijeess13.213>

Muhamet Kadrija^{1*}, Sabrije Kelmendi-Bardhaj², Besarta Pelaj³, Haxhi Kamberi^{4*}, NURSING CARE FOR PATIENTS AT THE END LIFE, page 109-124;

DOI: <https://doi.org/10.31407/ijeess13.214>

MUNTING DIVINE TII^{1*}, TCHOBSALA¹, LOCAL EXPLOITATION AND CHARACTERIZATION OF *BALANITES AEGYPTIACA* (L.) DEL PARKS IN THE SUDANO-SAHELIAN ZONE OF CAMEROON, FAR-NORTH, CAMEROON, page 125-138;

DOI: <https://doi.org/10.31407/ijeess13.215>

Ana T. Misoska¹, Snezhana Hristova^{1*}, Elena Bundaleska¹, Sanja Georgieva¹, HOW DO EMPLOYEES' ATTITUDES TOWARDS "GREEN BEHAVIOR" RELATE TO ENVIRONMENTAL CULTURE? page 139-152;

DOI: <https://doi.org/10.31407/ijeess13.216>

Ihor Ya. Tymochko¹, Oleksandr Yu. Chornobrov¹, Olha V. Bezrodnova², Ihor V. Solomakha¹, Oksana I. Drebot¹, Valentyna M. Maliarenko^{3*}, DEADWOOD STOCK IN A LINDEN-PINE-OAK FOREST OF SLOBOZHANSKYI NATIONAL NATURE PARK, page 153-162;

DOI: <https://doi.org/10.31407/ijeess13.217>

Hanna Myronova, Inna Honcharuk, Oleksandr Mazur, Oleksandr Tkachuk^{*}, Oksana Vradii, Olena Mazur, Yurii Shkatula, Liudmyla Peleh, Svitlana Okrushko, OPTIMIZATION OF MEASURES TO INCREASE DISEASE RESISTANCE OF POTATO VARIETIES AS A FACTOR OF REDUCING ENVIRONMENTAL POLLUTION, page 153-162;

DOI: <https://doi.org/10.31407/ijeess13.218>

OPTIMIZATION OF MEASURES TO INCREASE DISEASE RESISTANCE OF POTATO VARIETIES AS A FACTOR OF REDUCING ENVIRONMENTAL POLLUTION

Hanna Myronova, Inna Honcharuk, Oleksandr Mazur, Oleksandr Tkachuk*, Oksana Vradii, Olena Mazur, Yurii Shkatula, Liudmyla Peleh, Svitlana Okrushko

**Vinnytsia National Agrarian University, St. Sonyachna, 3, 21008, Vinnytsia, Ukraine;*

*Corresponding Author Oleksandr Tkachuk, email: tkachukop@ukr.net;

Received January 2022; Accepted February 2023; Published April 2023;

DOI: <https://doi.org/10.31407/ijeess13.218>

ABSTRACT

Potato varieties differed in their resistance to disease. The highest resistance to *Rhizoctonia violaceae* and Common scab was noted in the variety Granada. The affection of tubers of this variety by *Rhizoctonia violaceae* did not exceed 1.3%; the affection by Common scab did not exceed 1.8%. The higher affection of potato tubers by *Rhizoctonia violaceae* and Common scab was noted in the early-ripening variety Laperla and did not exceed 2.2 and 2.9%. The improvement of the agro background of cultivation provided a slight reduction in tuber affection – 0.2-0.3%, respectively. The variety Memphis was characterized by lower resistance to *Rhizoctonia violaceae* and Common scab compared to the variety Granada, but its resistance was higher compared to the variety Laperla. The affection of potato tubers by *Rhizoctonia violaceae* and Common scab in this variety did not exceed 1.8 and 2.8%. The affection of plants by viral diseases, providing the annual removal of diseased plants, depended primarily on the conditions of the year of research and varietal characteristics and varied from 0.5% in the variety Granada to 0.9% in the variety Memphis, while it did not exceed 0.7% in the variety Laperla. The conducted analysis of the influence of factors on disease affection indicated a significant share of the impact of genotype resistance – from 92.5 to 96.0%, as well as of the fraction of planting material – from 3.7 to 7.0%. The influence of other factors and their interaction was not significant. Therefore, the introduction of disease-resistant varieties into production and the use of the optimal fraction of planting material will reduce the pesticide load on the environment by reducing the volume of their use.

Key words: potato, diseases, stability.

INTRODUCTION

At the current stage of the development of global agricultural production, when its further intensification is taking place, such requirements as high and stable yield over the years, resistance to unfavourable growing conditions, diseases and pests, adaptability to mechanized cultivation, as well as high product quality are placed on the varieties. The variety, which is characterized by high yield potential in combination with reliable genetic protection of the crop against unfavourable environmental conditions, is the main factor in obtaining consistently high yields of potatoes (Kononuchenko et al., 2002; Biliavska et al., 2021; Mostovenko et al., 2022; Branitskyi et al., 2022).

Taking into account that potatoes in Ukraine are traditionally considered “second bread”, the use of high-quality planting material is the main prerequisite for highly productive potato growing. The share of seed material in crop growth is 25-30%, while it is more than 50% in unfavorable growing conditions. This is due to the fact that potatoes, which are propagated by tubers, gradually lose their seed qualities due to the accumulation of pathogens of various diseases in them (Vermenko et al., 2002).

Diseased plants have low viability, which causes a decrease in yield. Therefore, the main measure in obtaining high and stable yields of potatoes is the use of high-quality seed material (Bondarchuk, 2006).

MATERIALS AND METHODS

The research on the study of potato varieties was carried out at the farm “Olvia-S” in the village of Sopyn, Vinnytsia district, Vinnytsia region (Ukraine). The farm is located in the North-Eastern part of the Vinnytsia region.

The research was conducted during 2019-2021. The soil cover of the research area is represented by deep low-humus medium-loamy black soils. The basic seed potatoes were used for the research. Based on the working hypothesis and research objectives, we developed a three-factor experiment with four repetitions.

The air temperature during the growing season in 2019 was 17.1°C, which was by 0.8°C higher than the annual average data. In 2020 and 2021 the air temperature during the growing season was 16.4 and 16.5°C, which was as close as possible to the annual average indicators (Table 1).

Table 1. Characteristics of hydrothermal conditions during the research period

Month	Average monthly air temperature, °C				Amount of precipitation, mm			
	2019	2020	2021	Annual	2019	2020	2021	Annual average
April	9.3	9.2	9.3	9.1	37.8	32	26.8	48
May	15.4	11.7	13.7	14.9	144	136	116	54
June	21.6	20.2	18.9	18.1	88	68	98	83
July	19.0	20.3	19.5	20.2	38	30	54	91
August	20.2	20.4	21.0	19.4	9.2	28	28	69
In total, during the growing season	17.1	16.4	16.5	16.3	317	294	323	345

Regarding the amount of precipitation by month, there is a certain difference between the research years and the annual average data during the growing season. The largest amount of precipitation during the growing season (323 mm) was recorded in 2021, which was by 22 mm less than the annual average data, while it was by 6.0 mm and 29 mm more than the amount of precipitation in 2019 and 2020.

The air temperature during the years of field research shows that during 2019 there is an increase in the temperature regime compared to annual data. The conditions of 2021 turned out to be more favorable both in terms of moisture supply and temperature regime, which was reflected in the processes of growth and development of potatoes.

The basic seed potatoes were used for the research.

Based on the working hypothesis and research objectives, we developed a three-factor experiment with four repetitions.

- The area of the elementary plot is 38.5 m². Repetition is four times.
- Factor A – varieties: Laperla – early, Granada – mid-early, Memphis – mid-ripening.
- Factor B – the nutrition background and the method of applying mineral fertilizers. 40 t/ha of semi-rotted manure was applied under the predecessor (winter wheat). Mineral fertilizers such as Potassium magnesium (K₂₈Mg₈S₁₅) and Superphosphate (P₃₀) were applied for the main cultivation of potatoes. The complex mineral fertilizer Azofoska (N₁₅P₁₅K₁₅) was applied locally during planting or in pre-sowing cultivation.

- Factor C – fraction or mass of planting tubers: < 28 mm or 25-50 g; 28-60 mm (28-55 mm) or 51-80 g; > 60 (55) mm or 81-100 g. According to the experiment options in the varieties Laperla, Granada and Memphis, the costs of planting material on average were: 1) <28 mm or when planting tubers weighing 25-50 g – 2-2.15 t/ha; 2) 28-60 mm (28-55 mm) or when planting tubers weighing 51-80 g – 3.6-3.75 t/ha; 3) > 60 mm (55 mm) or when planting tubers weighing 81-100 g – 5.4-5.55 t/ha.

The field experiments were planned and carried out taking into account the requirements of the methodology of the experimental case [2, 3] and the methodological manual “Methodical recommendations for conducting research with potatoes” [2].

RESULTS AND DISCUSSION

During the period of research, a comparative assessment of disease affection of potato tubers was carried out, which mainly depended on the variety resistance to pathogens and the hydrothermal regime (Table 2).

Table 2. Disease affection and yield of potato varieties depending on fertilizer, fraction of planting tubers and varietal characteristics, 2019-2021

Fertilizer (factor B)	Fraction of planting tubers, mm (factor C)	Disease affection of tubers, %		Aukuba-mosaic	Yield, t/ha
		Rhizoctonia violaceae	Common scab		
Variety Laperla					
Without fertilizers (control)	1	2.2	2.9	0.7	19.5
	2	2.2	2.9	0.7	20.5
	3	2.1	2.8	0.7	21.4
40 t/ha of semi-rotted manure under the predecessor + K ₅₆ Mg ₁₆ S ₃₀ + P ₃₀ (background)	1	2.0	2.7	0.6	23.8
	2	2.0	2.7	0.6	25.0
	3	1.9	2.7	0.6	26.4
Background + N ₃₀ P ₃₀ K ₃₀ (local application)	1	1.9	2.6	0.6	27.2
	2	1.9	2.6	0.6	28.3
	3	1.9	2.6	0.6	29.9
Background + N ₄₅ P ₄₅ K ₄₅ (local application)	1	1.8	2.5	0.6	29.7
	2	1.8	2.5	0.6	31.1
	3	1.8	2.5	0.6	32.5
Background + N ₆₀ P ₆₀ K ₆₀ (scattering application)	1	1.9	2.6	0.6	28.0
	2	1.9	2.6	0.6	29.6
	3	1.9	2.6	0.6	31.1
Variety Granada					
Without fertilizers (control)	1	1.3	1.8	0.5	25.3
	2	1.3	1.8	0.5	26.3
	3	1.3	1.8	0.5	27.7
40 t/ha of semi-rotted manure under the predecessor + K ₅₆ Mg ₁₆ S ₃₀ + P ₃₀ (background)	1	1.2	1.7	0.5	31.6
	2	1.2	1.7	0.5	32.6
	3	1.1	1.6	0.5	34.3
Background +	1	1.1	1.6	0.4	36.3

N ₃₀ P ₃₀ K ₃₀ (local application)	2	1.1	1.6	0.4	37.6
	3	1.1	1.6	0.4	39.2
Background + N ₄₅ P ₄₅ K ₄₅ (local application)	1	1.0	1.5	0.4	38.3
	2	1.0	1.5	0.4	39.6
	3	1.0	1.5	0.4	41.1
Background + N ₆₀ P ₆₀ K ₆₀ (scattering application)	1	1.1	1.6	0.4	37.3
	2	1.1	1.6	0.4	38.5
	3	1.1	1.6	0.4	40.1
Variety Memphis					
Without fertilizers (control)	1	1.8	2.8	0.9	21.3
	2	1.8	2.8	0.9	22.1
	3	1.8	2.8	0.9	23.3
40 t/ha of semi-rotted manure under the predecessor + K ₅₆ Mg ₁₆ S ₃₀ +P ₃₀ (background)	1	1.7	2.7	0.8	30.0
	2	1.7	2.7	0.8	30.9
	3	1.7	2.7	0.8	32.6
Background + N ₃₀ P ₃₀ K ₃₀ (local application)	1	1.6	2.6	0.8	31.4
	2	1.6	2.6	0.8	32.9
	3	1.6	2.6	0.8	34.5
Background + N ₄₅ P ₄₅ K ₄₅ (local application)	1	1.5	2.5	0.8	33.2
	2	1.5	2.5	0.8	35.2
	3	1.5	2.5	0.8	36.7
Background + N ₆₀ P ₆₀ K ₆₀ (scattering application)	1	1.6	2.6	0.8	32.9
	2	1.6	2.6	0.8	34.0
	3	1.6	2.6	0.8	35.6
<i>HIP₀₅: 2019 A-0.16; B-0.16; C-0.21; AB-0.29; AC-0.36; BC-0.37; ABC-0.64;</i> <i>2020 A-0.15; B-0.15; C-0.2; AB-0.27; AC-0.35; BC-0.36; ABC-0.62;</i> <i>2021 A-0.24; B-0.24; C-0.31; AB-0.42; AC-0.53; BC-0.53; ABC-0.93</i>					

It should be noted that potato varieties differed in their resistance to disease. The highest resistance to *Rhizoctonia violaceae* and Common scab was noted in the variety Granada. The affection of tubers of this variety by *Rhizoctonia violaceae* did not exceed 1.3%; the affection by Common scab did not exceed 1.8%.

In general, the affection of tubers by both *Rhizoctonia violaceae* and Common scab was reduced by 0.1-0.3% with the intensification of technological methods of cultivation (application of different methods and rates of fertilizers, fractions of planting material).

The analysis of influencing factors on the studied elements shows that the predominant share of the impact on the affection of potato varieties by *Rhizoctonia violaceae* belongs to genotypic characteristics – 92.48%. The impact of the fraction of planting material was also significant – 7.0%. The share of the influence of fertilizer, as well as the interaction of these factors, was lower, except for the interaction of the fraction of planting material with fertilizer – 0.3%, which turned out to be significant (Fig. 1).

The higher affection of potato tubers by *Rhizoctonia violaceae* and Common scab was noted in the early-ripening variety Laperla, but it did not exceed 2.2 and 2.9%. The improvement of the agro background of cultivation provided a slight reduction in tuber affection – 0.2-0.3%, respectively.

The variety Memphis was characterized by lower resistance to *Rhizoctonia violaceae* and Common scab compared to the variety Granada, but its resistance was higher compared to the variety Laperla. The affection of potato tubers by *Rhizoctonia violaceae* and Common scab in this variety did not exceed 1.8 and 2.8%. The use of the researched technological methods of cultivation contributed to a slight decrease in tuber affection – 0.2-0.3%. A slight increase in affection by *Rhizoctonia violaceae* was noted in 2020.

It should be noted that the share of influence on the affection by Common scab mostly depended on genotypic characteristics (96.0%) and on the fraction of planting material (3.7%). The share of the influence of fertilizer, as well as the interaction of these factors, was lower (Fig. 2).

The plant affection by viral diseases was determined in field conditions in the phase of budding and flowering. At the same time, the affection of plants by viral diseases, providing the annual removal of diseased plants, depended primarily on the conditions of the year of research and varietal characteristics and varied from 0.5% in the variety Granada to 0.9% in the variety Memphis, while it did not exceed 0.7% in the variety Laperla. A slightly higher affection by viral diseases was noted in 2019 and 2020. Since the year 2019 was hotter than others, and the year 2020 saw a slightly lower amount of precipitation, that is, these years turned out to be more favorable for the flight of aphids.




Figure 1. Shares of influence of varietal characteristics, fractions of planting material and fertilizer on the affection of potato varieties by *Rhizoctonia violaceae*

The affection by Aukuba-mosaic depended on genotypic characteristics (95.05%) and on the fraction of planting material (4.33%). The share of influence of fertilizer, as well as the interaction of these factors, was lower (Fig. 3).




Figure 2. Shares of influence of varietal characteristics, fractions of planting material and fertilizer on the affection of potato varieties by Common scab.

The yield of potatoes depended on the methods and rates of fertilizers, fraction of planting material and varietal characteristics. It should be noted that an increase in the fraction of planting tubers led to an increase in the level of potato yield under different methods and rates of applied fertilizers among all potato varieties studied in the experiments.

Among the studied varieties, the yield indicators of the variety Granada were the best, which was noted in all options of the experiment. The highest productivity indicators were noted in the experimental option, where local application of mineral fertilizers in the dose of $N_{45}P_{45}K_{45}$ was carried out against the background of the effect of semi-rotted manure applied under the predecessor.

The yield increased from 38.3 to 41.1 t/ha. It is by 13.0-13.4 t/ha higher than in the control option. The experimental option, where mineral fertilizers in the dose of $N_{60}P_{60}K_{60}$ were applied by scattering against the background of Potassium magnesium fertilizer and the effect of semi-rotted manure, provided a lower yield from 37.3 to 40.1 t/ha, compared to the previous option.

The fraction of planting material increased. It is by 12.0-12.4 t/ha higher than in the control option and indicates a lower efficiency of the effect of scattering application compared to local application, during which fertilizers are placed at a certain soil depth with better moisture supply. A zone with an increased concentration of nutrients is formed, which are more fully used by plants during the growing season.




Figure 3. Shares of influence of varietal characteristics, fractions of planting material and fertilizer on the affection of potato varieties by Aukuba-mosaic.

The mid-ripening variety Memphis, according to all experimental options, provided intermediate indicators in terms of the elements of the crop structure and yield level between the mid-early variety Granada and the early-ripening variety Laperla, which are better in terms of these indicators.

CONCLUSIONS

- Potato varieties differed in their resistance to disease. The highest resistance to *Rhizoctonia violaceae* and Common scab was noted in the variety Granada. The affection of tubers of this variety by *Rhizoctonia violaceae* did not exceed 1.3%; the affection by Common scab did not exceed 1.8%. The higher affection of potato tubers by *Rhizoctonia violaceae* and Common scab was noted in the early-ripening variety Laperla and did not exceed 2.2 and 2.9%. The improvement of the agro background of cultivation provided a slight reduction in tuber affection – 0.2-0.3%, respectively. The variety Memphis was characterized by lower resistance to *Rhizoctonia violaceae* and Common scab compared to the variety Granada, but its resistance was higher compared to the variety Laperla. The affection of potato tubers by *Rhizoctonia violaceae* and Common scab in this variety did not exceed 1.8 and 2.8%. The affection of plants by viral diseases, providing the annual removal of diseased plants, depended primarily on the conditions of the year of research and varietal characteristics and varied from 0.5% in the variety Granada to 0.9% in the variety Memphis, while it did not exceed 0.7% in the variety Laperla.

- The conducted analysis of the influence of factors on disease affection indicated a significant share of the impact of genotype resistance – from 92.5 to 96.0%, as well as of the fraction of planting material – from 3.7 to 7.0%. The influence of other factors and their interaction was not significant. Therefore, the introduction of disease-resistant varieties into production and the use of the optimal fraction of planting material will reduce the pesticide load on the environment by reducing the volume of their use.

REFERENCES

1. Biliavska L., Biliavskiy Y., Mazur O., Mazur O. (2021) Adaptability and breeding value of soybean varieties of Poltava breeding. Bulgarian Journal of Agricultural Science. 27 (2), 312-322;
2. Bondarchuk A. A. (2006) Seed production is the main factor of high-yield potato cultivation. Potato production of Ukraine. 1-2, 7-8;
3. Branitskyi Yu., Telekalo N., Kupchuk I., Mazur O., Alieksieiev O., Okhota Yu., Mazur O. (2021) Improvement of technological methods of switchgrass (*Panicum virgatum* L.) growing in the Vinnytsia region. Acta phytotechn zootech. 25 (4), 311-318;
4. Didur I., Bakhmat M., Chynchyk O., Pansyryeva H., Telekalo N., Tkachuk O. (2020) Substantiation of agroecological factors on soybean agrophytocenoses by analysis of variance of the Right-Bank Forest-Steppe in Ukraine. Ukrainian Journal of Ecology. 10 (5), 54-61;
5. Didur I., Chynchyk O., Pansyryeva H., Olifirovych S., Olifirovych V., Tkachuk O. (2021) Effect of fertilizers for *Phaseolus vulgaris* L. productivity in Western Forest-Steppe of Ukraine. Ukrainian Journal of Ecology. 11 (1), 419-424;
6. Ermantraut E.R., Prysiazhniuk O.I., Shevchenko L.I. (2007) Statistical analysis of agronomic research data in the STATISTICA 6.0 package. Kyiv: Poligraf Consulting. 55;
7. Honcharuk I., Tkachuk O., Mazur O., Kravets R., Mazur R., Alieksieiev O., Tetyana Z., Bronnikova L. (2023) Manifestation of ecological-adaptive properties of soybean varieties depending on soil-climatic conditions. International Journal of Ecosystems and Ecology Science. 13, 2, 51-60;
8. Kononuchenko V.V., Molotskyi M.Ya. (2002) Potato. Bila Tserkva. 1, 536;
9. Kutsenko V.S., Osypchuk A.A., Podgaetskyi A.A. (2002) Methodological recommendations for conducting research with potatoes. Nemishaeve: Intas. 182;
10. Mazur V., Didur I., Tkachuk O., Pansyryeva H., Ovcharuk V. (2021) Agroecological stability of cultivars of sparsely distributed legumes in the context of climate change. Scientific Horizons. 24 (1), 54-60;
11. Mazur V., Tkachuk O., Pansyryeva H., Demchuk O. (2021) Quality of pea seeds and agroecological condition of soil when using structured water. Scientific Horizons. 24 (7), 53-60;
12. Mazur V., Tkachuk O., Pansyryeva H., Kupchuk I., Mordvaniuk M., Chynchyk O. (2021) Ecological suitability peas (*Pisum sativum*) varieties to climate change in Ukraine. Agraarteadus. Journal of Agricultural Science. 2 (32), 276-283;
13. Mostovenko V., Mazur O., Didur I., Kupchuk I., Voloshyna O., Mazur O. (2022) Garden pea yield and its quality indicators depending on the technological methods of growing in conditions of Vinnytsia region. Acta phytotechn zootech. 25 (3), 226-241;
14. Razanov S., Husak O., Polishchuk M., Bakhmat O., Koruniak O., Symochko L., Ovcharuk I. (2022) Accumulation peculiarities of heavy metals in cereal crops grains of different vegetation period in conditions of the forest steppe of the right bank of Ukraine. International Journal of Ecosystems and Ecology Science (IJEES). 12 (3), 43-50. <https://doi.org/10.31407/ijeess12.3>;
15. Razanov S., Piddubna A., Gucol G., Symochko L., Kovalova S., Bakhmat M., Bakhmat O. (2022) Estimation of heavy metals accumulation by vegetables in agroecosystems as one of the main aspects in food security. International Journal of Ecosystems and Ecology Science (IJEES). 12 (3), 159-164. <https://doi.org/10.31407/ijeess12.320>;
16. Razanov S.F., Tkachuk O.P., Bakhmat O.M., Razanova A.M. (2020). Reducing danger of heavy metals accumulation in winter wheat grain which is grown after leguminous perennial precursor. Ukrainian Journal of Ecology. 10 (1), 254-260. doi: 10.15421/2020_40;
17. Razanov S.F., Tkachuk O.P., Razanova A.M., Bakhmat M.I., Bakhmat O.M. (2020) Intensity of heavy metal accumulation in plants of *Silybum marianum* L. in conditions of field rotation. Ukrainian Journal of Ecology. 10 (2), 131-136. doi: 10.15421/2020_75;

18. Razanov S.F., Tkachuk O.P., Mazur V.A., Didur I.M. (2018) Effect of bean perennial plants growing on soil heavy metal concentrations. *Ukrainian Journal of Ecology*. 8 (2), 294-300. doi: 10.15421/2018_341;
19. Tkachuk O., Telekalo N. (2020) Agroecological potential of legumes in conditions of intensive agriculture of Ukraine. Collective monograph. Publishing House «Baltija publishing». Riga. Latvia. 42-108;
20. Tkachuk O., Verhelis V. (2021) Intensity of soil pollution by toxic substances depending on the degree of its washout. *Scientific Horizons*. 24 (3), 52-57;
21. Tkachuk O. (2021) Biological features of the distribution of root systems of perennial legume grasses in the context of climate change. *Scientific Horizons*. 24 (2), 70-76;
22. Tkachuk O., Mordvaniuk M. (2021) Research of technological indicators of grain quality of legumes as objects of storage and processing. Development of scientific, technological and innovation space in Ukraine and EU countries. Riga, Latvia: «Baltija Publishing». 221-240;
23. Tkachuk O., Mordvanyuk M. (2022) Study of the influence of unfavorable vegetation conditions on agro-ecological resistance of bean varieties. Theoretical and practical aspects of the development of modern scientific research: Scientific monograph. Publishing House «Baltija Publishing», Riga, Latvia. 109-125;
24. Vermenko Yu.Ya., Molotskyi M.Ya. (2002) Causes of potato degeneration and measures to combat them. *Potato*.1, 379-395;